

# Comment on: ‘Universal Behavior of Load Distribution in Scale-free Networks’

In a previous Letter [1], Goh *et al* have presented a numerical study of the load—or betweenness centrality—distribution in a scale-free network whose degree distribution follows a power law  $p(k) \sim k^{-\gamma}$  where  $\gamma \in [2, \infty[$  is a tunable parameter. They showed that the load  $\ell$  is distributed according to a power-law  $P(\ell) \sim \ell^{-\delta}$  with exponent  $\delta$ . On the basis of their numerical results, they conjectured that the value of  $\delta \simeq 2.2$  is independent of  $\gamma$  for the interval  $[2, 3)$ . Based on this apparent universality, a classification of scale-free networks according to the value of  $\delta \simeq 2.2$  (class I) or  $\delta = 2$  (class II) was proposed [2]. In this comment we argue that the value of  $\delta$  is not universal and varies significantly as  $\gamma$  changes in the interval  $[2, 3)$ . The power law fits of the cumulative function  $\text{Prob}(\text{load} \geq \ell)$  for the model proposed in [1] gives the values  $\delta = 1.84 \pm 0.04$ ,  $2.05 \pm 0.05$  and  $2.25 \pm 0.05$  for  $\gamma = 2, 2.5$ , and  $3$  respectively, while for the BA model [3]  $\delta = 2.3 \pm 0.1$ . The variations of  $\delta$  are significant enough to claim that it is not universal but in order to double-check our results we use an indirect way of computing  $\delta$ . We study the relation between the load and the connectivity [1,4] which is of the form  $\ell \sim k^\eta$  where the exponent  $\eta$  depends on the network. As can be seen on Fig. (1a), the power law holds remarkably for a large range of  $k$  allowing for an accurate measure of  $\eta$ . We also checked that the value of  $\eta$  does not change significantly for different values of the system size (For  $\gamma = 2.5$ , we obtain a relative variation due to size going from  $N = 10^4$  to  $5 \cdot 10^4$  less than 1%). The exponents  $\eta$  and  $\delta$  are not independent and it is easy to show that [4]  $\eta = (\gamma - 1)/(\delta - 1)$ . If the value of  $\delta \simeq 2.2$  is universal then  $\eta$  is a linear function of  $\gamma$  with slope  $\simeq 1/1.2 \simeq 0.83$ . In Fig. (1b) we plot the measured  $\eta$  versus  $\gamma$  for the different types of networks studied and the corresponding value predicted by universality. This Fig. (1b) shows that if for  $\gamma \simeq 3$  the value  $\delta = 2.2$  seems to be acceptable, the claim of universality for  $\gamma \in [2, 3)$  proposed in [1] does not hold (our results do not fit in the other class  $\delta = 2.0$  either). In addition, we tested the universality for different values of  $m$  and we also obtain variations ruling it out: For  $\gamma = 2.5$  and for  $N = 2 \cdot 10^4$ , we obtain  $\eta = 1.477 \pm 0.006$ ,  $1.56 \pm 0.006$ , and  $1.64 \pm 0.01$  for  $m = 2, 4, 6$  respectively.




FIG. 1. (a) Log-Log plot of the normalized average load versus connectivity for the same models as in [1] with  $m = 2$ . The power law fits (straight lines) give  $\eta = 1.27 \pm 0.01$  ( $N = 3 \cdot 10^4$ ),  $1.467 \pm 0.006$  ( $N = 5 \cdot 10^4$ ), and  $1.68 \pm 0.02$  ( $N = 5 \cdot 10^4$ ) for  $\gamma = 2, 2.5$ , and  $3$  respectively. For the BA model,  $\eta = 1.81 \pm 0.02$  ( $N = 5 \cdot 10^4$ ). (b)  $\eta$  versus  $\gamma$ . If the universality proposed in [1] would be correct, the measured values for  $\gamma \in [2, 3)$  should lie on the “universal” straight line corresponding to  $\delta = 2.2$  (class I).

The important exponent thus appears to be  $\eta$  and it is interesting to note that  $\eta$  is significantly smaller than the maximum value  $\eta = 2$ . This maximum value is reached when nodes with large centrality (ie. with large  $k$ ) link together disconnected parts of roughly the same size. The load of these nodes is then of the order of  $\ell \sim k(k-1)/2 \sim k^2$ . The fact that  $\eta < 2$  indicates that the different parts are also connected by shortest paths which do not pass through the central node. More generally, it would be interesting to understand how  $\eta$  depends on the different parameters of the network such as  $\gamma$  and the degree correlation.

Marc Barthélemy

CEA-Service de Physique de la Matière Condensée  
BP12 Bruyères-Le-Châtel, France

- 
- [1] K.-I. Goh, B. Kahng, and D. Kim, Phys. Rev. Lett. **87**, 278701 (2001).
  - [2] K.-I. Goh, H. Jeong, B. Kahng, and D. Kim, Proc. Natl. Acad. Sci. (USA) **99**, 12583 (2002).
  - [3] A.-L. Barabasi and R. Albert, Science **286**, 509 (1999).
  - [4] A. Vazquez, R. Pastor-Satorras, and A. Vespignani, Phys. Rev. E **65**, 066130 (2002).